

A COMPLETELY INTEGRABLE SYSTEM ON G_2 COADJOINT ORBITS

JEREMY LANE

ABSTRACT. We construct a Gelfand-Zeitlin system on a one-parameter family of G_2 coadjoint orbits that are multiplicity-free Hamiltonian $SU(3)$ -spaces. Using this system we prove a lower bound for the Gromov width of these orbits. This lower bound agrees with the known upper bound.

1. INTRODUCTION

Gelfand-Zeitlin systems are completely integrable systems constructed from maximal non-Abelian Hamiltonian symmetries of a symplectic manifold. The classic examples of Gelfand-Zeitlin systems were constructed on $SU(n)$ and $SO(n)$ coadjoint orbits [6]. These systems were employed by [16] to prove tight lower bounds for the Gromov width of $SU(n)$ and $SO(n)$ coadjoint orbits that, together with the upper bounds obtained in [4], compute the Gromov width. Tight lower bounds were also recently proven for $Sp(n)$ coadjoint orbits using integrable systems constructed from toric degenerations [8].

Motivated by the work of [16], and a desire for more examples of Gelfand-Zeitlin systems, this paper describes a Gelfand-Zeitlin system on a one-parameter family of G_2 coadjoint orbits that are multiplicity-free $SU(3)$ -spaces. As complex manifolds, these coadjoint orbits are degree 18, 5 dimensional projective subvarieties of \mathbb{CP}^{13} [1]. Using the convexity theorem of [13] we are able to compute the image of the Gelfand-Zeitlin system – a 5-dimensional convex polytope – without undue difficulty and this image classifies an open dense submanifold of the coadjoint orbit up to symplectomorphism. Combining the lower bound methods employed recently in [16], and the upper bounds due to [4], we can then prove

Theorem 1. *If \mathcal{O}_λ is the G_2 coadjoint orbit through $(0, \lambda)$ as in Figure 1 equipped with the Kostant-Kirillov-Souriau symplectic form ω_λ , then*

$$GWidth(\mathcal{O}_\lambda, \omega_\lambda) = \frac{\lambda}{\sqrt{3}}.$$

The other one-parameter family of 5 (complex) dimensional G_2 coadjoint orbits, corresponding to the short simple root, are quadrics in \mathbb{CP}^6 that are homogeneous spaces for $SO(7, \mathbb{C})$. As such, they are $SO(8)$ coadjoint orbits and carry Gelfand-Zeitlin systems already described in [16] that, combined with the upper bound of [4], give the precise Gromov width of these orbits.

One may also deduce the Gromov width of both these families of 5 dimensional G_2 coadjoint orbits from the recent work of [14], since they have second Betti number $b_2 = 1$. The lower bound in the main theorem of *op. cit.* follows from work of [9, 12] that constructs completely integrable torus actions on projective varieties from toric degenerations. In comparison, the virtue of this work is that the momentum map and polytope have a very straightforward descriptions.

2010 *Mathematics Subject Classification.* 37J35, 57R17 (primary), 53D20 (secondary). Please refer to <http://www.ams.org/msc/> for a list of codes.

Acknowledgements The author would like to thank his advisor Yael Karshon as well as Milena Pabiniak and Alexander Caviedes Castro for sharing their mathematical insights. This work was supported by a NSERC PGS scholarship.




FIGURE 1

2. GELFAND-ZEITLIN SYSTEMS AND MULTIPLICITY-FREE SPACES

Introduced by [6], Gelfand-Zeitlin systems can be viewed as examples of the following general construction, a detailed exposition of which can be found in [13].

Let G be a compact, connected Lie group and let (M, ω, Φ, G) be a connected Hamiltonian G -space. Given a chain of subgroups $H_1 \leq \dots \leq H_n = G$ with corresponding subalgebras $\mathfrak{h}_k = \text{Lie}(H_k)$, there are dual projection maps $p_k : \mathfrak{h}_{k+1}^* \rightarrow \mathfrak{h}_k^*$. Fixing closed positive Weyl chambers, $\mathfrak{t}_{k,+}^* \subset \mathfrak{h}_k^*$, we can construct the map

$$(2) \quad \begin{aligned} \Lambda : \mathfrak{g}^* &\rightarrow \mathcal{T} = \mathfrak{t}_{n,+}^* \times \dots \times \mathfrak{t}_{1,+}^* \subset \mathfrak{t}_n^* \times \dots \times \mathfrak{t}_1^* \\ \Lambda &= (s, s \circ p_{n-1}, \dots, s \circ p_1 \circ \dots \circ p_{n-1}). \end{aligned}$$

where $s(\xi)$ is defined as the unique element of $(H_k \cdot \xi) \cap \mathfrak{t}_{k,+}^*$. The restriction of the map

$$(3) \quad \Lambda \circ \Phi : M \rightarrow \mathcal{T}$$

to the open, dense submanifold $U = (\Lambda \circ \Phi)^{-1}(\mathcal{T}^{\text{int}})$ is a smooth momentum map for a Hamiltonian torus action¹. This construction is referred to as *Thimm's trick* or *the method of Thimm*, and one may say that the map $\Lambda \circ \Phi$ is *constructed by Thimm's trick*, in reference to the paper [18].

Recall that a Hamiltonian G -space is *multiplicity-free* if for all $\xi \in \mathfrak{g}^*$ the reduced space $\Phi^{-1}(\xi)/G_\xi$ is a point. An effective Hamiltonian torus action is completely integrable if the dimension of the torus is half the real dimension of the manifold. If M is a multiplicity-free G -space, and for each $1 \leq k < n$, every H_{k+1} coadjoint orbit is a multiplicity-free H_k -space, the Hamiltonian torus action induced on U is completely integrable [7]. Following Guillemin and Sternberg's terminology in [6], we refer to such a map (and the associated densely defined torus action) as a *Gelfand-Zeitlin² system*.

¹To be more precise, one should replace each of the closed chambers $\mathfrak{t}_{k,+}^*$ in (2) with the closure of the principal face of $\mathfrak{t}_{k,+}^*$ corresponding to the induced Hamiltonian H_k -actions on M .

²One should note that there are multiple spellings of Zeitlin in the literature, including Tsetlin and Cetlin. Works by Kostant-Wallach, Kogan-Miller, and Guillemin-Sternberg respectively each choose a different spelling.

Example 4. Let \mathcal{O} be a $U(k)$ coadjoint orbit, and define the subgroup

$$H_{k-1} = \{\text{diag}(A, 1) : A \in U(k-1)\} \cong U(k-1).$$

Guillemin and Sternberg showed that \mathcal{O} is a multiplicity-free $U(k-1)$ -space. Thus if M is a multiplicity-free $SU(n)$ -space, then the map $\Lambda \circ \Phi$ constructed on M by Thimm's trick for the chain of subgroups

$$U(1) \leq U(2) \leq \cdots \leq U(n-1) \leq SU(n)$$

is a Gelfand-Zeitlin system.

If $\Lambda \circ \Phi : M \rightarrow \mathcal{T}$ is a map constructed by Thimm's trick, then the submanifold U is connected. Using this fact and the convexity theorem for proper torus momentum maps [2, 3] one can prove

Theorem 5. [13] *Let M be a connected symplectic manifold and suppose $\Lambda \circ \Phi : M \rightarrow \mathcal{T}$ is a map constructed by Thimm's trick. If $\Lambda \circ \Phi$ is a proper map, then $\Lambda \circ \Phi(M)$ is convex and the fibres of $\Lambda \circ \Phi$ are connected.*

3. A GELFAND-ZEITLIN SYSTEM ON \mathcal{O}_λ

In [20], Chris Woodward gives the following example.

Example 6. Consider the exceptional Lie group G_2 with maximal torus T and let \mathfrak{t} be the corresponding subalgebra. Fix the standard identifications $\mathfrak{t} \cong \mathfrak{t}^* \cong \mathbb{R}^2$ and let

$$S = \left\{ \alpha_1 = \left(\frac{-3}{2}, \frac{\sqrt{3}}{2} \right), \alpha_2 = (1, 0) \right\} \subset \mathbb{R}^2$$

be the standard choice of simple roots (see Figure 1A). The positive Weyl chamber $\mathfrak{t}_{G_2,+}^*$ given by this choice of simple roots is shaded in Figure 1B. There is an embedding of $SU(3)$ in G_2 with maximal torus T corresponding to the long roots of G_2 , as shown in Figure 2A.

Coadjoint G_2 -orbits are parameterized by points in $\mathfrak{t}_{G_2,+}^*$. In particular, we write \mathcal{O}_λ for the G_2 coadjoint orbit through the point $(0, \lambda)$ on the vertical edge of $\mathfrak{t}_{G_2,+}^*$, equipped with its Kostant-Kirillov-Souriau symplectic form ω_λ . The following facts are explained in [20].

- (1) The coadjoint orbit $\mathcal{O}_\lambda \cong G_2/(SU(2) \times U(1))$ has real dimension 10.
- (2) The $SU(3)$ subgroup acts locally freely on \mathcal{O}_λ . Since $\dim SU(3) + \text{rank} SU(3) = 10$, \mathcal{O}_λ is a multiplicity free $SU(3)$ -space.
- (3) The Kirwan polytope for the $SU(3)$ -action can be understood using the Heckman formula of [5] and is the triangle illustrated in Figure 2B. In coordinates, the three vertices of this triangle are

$$(0, \lambda), \left(\frac{\lambda}{2\sqrt{3}}, \frac{\lambda}{2} \right), \left(\frac{-\lambda}{2\sqrt{3}}, \frac{\lambda}{2} \right).$$

In terms of the roots of G_2 the vertices are

$$\frac{\lambda(\alpha_4 - \alpha_5)}{\sqrt{3}}, \frac{-\lambda\alpha_5}{\sqrt{3}}, \frac{\lambda\alpha_4}{\sqrt{3}}.$$

Following the previous section, consider the chain of groups

$$U(1) \leq U(2) \leq SU(3)$$

where we choose the embeddings

$$U(1) \cong \{\text{diag}(a, 1) : a \in U(1)\} \leq U(2), U(2) \cong \left\{ \frac{1}{\det(A)} \text{diag}(A, 1) : A \in U(2) \right\} \leq SU(3).$$




FIGURE 2

Since \mathcal{O}_λ is a multiplicity-free $SU(3)$ -space, the map

$$(7) \quad \Lambda \circ \Phi : \mathcal{O}_\lambda \rightarrow \mathfrak{t}_{SU(3),+}^* \times \mathfrak{t}_{U(2),+}^* \times \mathfrak{t}_{U(1),+}^*$$

is a Gelfand-Zeitlin system. For the remainder of this paper $\Lambda \circ \Phi$ will denote this specific Gelfand-Zeitlin system.

4. THE GELFAND-ZEITLIN POLYTOPE OF \mathcal{O}_λ

In this section we compute the image of the Gelfand-Zeitlin system on \mathcal{O}_λ . In order to do this we compute the vertices of the image, a list of bounding inequalities, and apply Theorem 5.

Fix the standard identifications

$$\begin{aligned} \mathfrak{u}(n) &\cong \{X \in M_n \mathbb{C} : X = \overline{X}^t\} \\ \mathfrak{t}_{U(n)}^* &\cong \mathfrak{t}_{U(n)} \cong \{\text{diag}(x_1, \dots, x_n) : x_1, \dots, x_n \in \mathbb{R}\} \cong \mathbb{R}^n \\ L_{U(n)}^* &= \text{Hom}(L_{U(n)}, \mathbb{Z}) \cong L_{U(n)} \cong \{\text{diag}(x_1, \dots, x_n) : x_1, \dots, x_n \in \mathbb{Z}\} \cong \mathbb{Z}^n \\ \mathfrak{t}_{U(n),+}^* &\cong \{\text{diag}(x_1, \dots, x_n) \in \mathfrak{t}_u(n) : x_1 \leq \dots \leq x_n\} \end{aligned}$$

From the previous section we also have the identifications

$$\begin{aligned} \mathfrak{t}_{SU(3)}^* &= \mathbb{R}^2 \\ L_{SU(3)}^* &= \mathbb{Z} \left\langle \alpha_2 = (1, 0), \alpha_4 = \left(\frac{-1}{2}, \frac{\sqrt{3}}{2} \right) \right\rangle. \end{aligned}$$

With these identifications the Gelfand-Zeitlin system (7) is a map to $\mathcal{T} \subset \mathbb{R}^5 = \mathbb{R}^2 \times \mathbb{R}^2 \times \mathbb{R}^1 \cong \mathfrak{t}_{SU(3)}^* \times \mathfrak{t}_{U(2)}^* \times \mathfrak{t}_{U(1)}^*$ with coordinates $(x_1, x_2, x_3, x_4, x_5)$, where

$$\begin{aligned} \mathcal{T} &= \left\{ (x_1, x_2, x_3, x_4, x_5) \in \mathbb{R}^5 : \frac{\pm 3x_1}{2} + \frac{\sqrt{3}x_2}{2} \geq 0 \text{ and } x_3 \leq x_4 \right\} \\ &\cong \mathfrak{t}_{SU(3),+}^* \times \mathfrak{t}_{U(2),+}^* \times \mathfrak{t}_{U(1),+}^* \end{aligned}$$

Further, in these coordinates the weight lattice $L^* = \text{Hom}(L, \mathbb{Z})$ is

$$\begin{aligned} (8) \quad L^* &= L_{SU(3)}^* \times L_{U(2)}^* \times L_{U(1)}^* \\ &= \mathbb{Z} \left\langle (1, 0, 0, 0, 0), \left(\frac{1}{2}, \frac{\sqrt{3}}{2}, 0, 0, 0 \right), (0, 0, 1, 0, 0), (0, 0, 0, 1, 0), (0, 0, 0, 0, 1) \right\rangle. \end{aligned}$$

TABLE 1. Vertices of the Gelfand-Zeitlin polytope Δ_λ

$$\begin{array}{c}
(x_1, x_2, x_3, x_4, x_5) \\
\hline
(0, \lambda, 0, 0, 0) \\
\left(0, \lambda, \frac{-\lambda}{\sqrt{3}}, 0, \frac{-\lambda}{\sqrt{3}}\right) \\
\left(0, \lambda, \frac{-\lambda}{\sqrt{3}}, \frac{\lambda}{\sqrt{3}}, \frac{-\lambda}{\sqrt{3}}\right) \\
\left(0, \lambda, 0, \frac{\lambda}{\sqrt{3}}, 0\right) \\
\left(0, \lambda, 0, \frac{\lambda}{\sqrt{3}}, \frac{\lambda}{\sqrt{3}}\right) \\
\left(0, \lambda, \frac{-\lambda}{\sqrt{3}}, \frac{\lambda}{\sqrt{3}}, \frac{\lambda}{\sqrt{3}}\right) \\
\left(0, \lambda, \frac{-\lambda}{\sqrt{3}}, 0, 0\right) \\
\left(\frac{-2\lambda}{3\sqrt{3}}, \frac{\lambda}{3\sqrt{3}}, \frac{-2\lambda}{3\sqrt{3}}, \frac{-2\lambda}{3\sqrt{3}}, \frac{-2\lambda}{3\sqrt{3}}\right) \\
\left(\frac{-2\lambda}{3\sqrt{3}}, \frac{\lambda}{3\sqrt{3}}, \frac{-2\lambda}{3\sqrt{3}}, \frac{\lambda}{3\sqrt{3}}, \frac{\lambda}{3\sqrt{3}}\right) \\
\left(\frac{-2\lambda}{3\sqrt{3}}, \frac{\lambda}{3\sqrt{3}}, \frac{-2\lambda}{3\sqrt{3}}, \frac{\lambda}{3\sqrt{3}}, \frac{-2\lambda}{3\sqrt{3}}\right) \\
\left(\frac{-\lambda}{3\sqrt{3}}, \frac{2\lambda}{3\sqrt{3}}, \frac{-\lambda}{3\sqrt{3}}, \frac{-\lambda}{3\sqrt{3}}, \frac{-\lambda}{3\sqrt{3}}\right) \\
\left(\frac{-\lambda}{3\sqrt{3}}, \frac{2\lambda}{3\sqrt{3}}, \frac{-\lambda}{3\sqrt{3}}, \frac{2\lambda}{3\sqrt{3}}, \frac{-\lambda}{3\sqrt{3}}\right) \\
\left(\frac{-\lambda}{3\sqrt{3}}, \frac{2\lambda}{3\sqrt{3}}, \frac{-\lambda}{3\sqrt{3}}, \frac{2\lambda}{3\sqrt{3}}, \frac{2\lambda}{3\sqrt{3}}\right)
\end{array}$$

The Kirwan polytope of the previous section gives us inequalities

$$\begin{aligned}
(9) \quad & \frac{-\sqrt{3}x_1}{2} + \frac{x_2}{2} \leq \frac{\lambda}{2}, \\
& \frac{\sqrt{3}x_1}{2} + \frac{x_2}{2} \leq \frac{\lambda}{2}, \\
& \frac{\lambda}{2} \leq x_2.
\end{aligned}$$

From the interlacing inequalities for eigenvalues of Hermitian matrices, one derives

$$\begin{aligned}
(10) \quad & \frac{-x_1}{3} - \frac{x_2}{\sqrt{3}} \leq x_3 \leq \frac{2x_1}{3} \leq x_4 \leq \frac{-x_1}{3} + \frac{x_2}{\sqrt{3}} \\
& x_3 \leq x_5 \leq x_4.
\end{aligned}$$

One can find elements of $\Phi(\mathcal{O}_\lambda)$ that make each of the inequalities strict, so the image of $\Lambda \circ \Phi$ contains the points listed in Table 1. By Theorem 5 and the inequalities (9) and (10) we have

Proposition 11. *The image of the Gelfand-Zeitlin system $\Lambda \circ \Phi : \mathcal{O}_\lambda \rightarrow \mathbb{R}^5$ is the convex hull of the 13 points listed in Table 1. This convex hull, which we will refer to as Δ_λ , is a 5-dimensional convex polytope.*

5. THE GROMOV WIDTH OF \mathcal{O}_λ

The *Gromov width* of a symplectic manifold is defined as

$$\text{GWidth}(M, \omega) = \sup \{ \pi r^2 > 0 : \exists \text{ a symplectic embedding } B^{2n}(r) \rightarrow M \}$$

where $B^{2n}(r)$ is the open ball of radius r in \mathbb{R}^{2n} equipped with the standard symplectic structure. Coadjoint orbits have the Kostant-Kirillov-Souriau symplectic structure and

their Gromov width has been a subject of much study, including work on complex Grassmannians by [11, 15] and later other families of coadjoint orbits [21, 16, 4, 8, 14]. In this section we follow the same approach as [16] to compute a lower bound for the Gromov width of \mathcal{O}_λ from the polytope Δ_λ .

Suppose a connected symplectic manifold (M, ω) is equipped with a completely integrable action of a torus T generated by a momentum map

$$\mu : M \rightarrow \mathfrak{t}^*.$$

If there is an open, convex set $\mathcal{C} \subset \mathfrak{t}^*$ such that $\mu(M) \subset \mathcal{C}$ and the map $\mu : M \rightarrow \mathcal{C}$ is proper, then M is classified up to T -equivariant symplectomorphism by its image, $\mu(M)$ ([10], Proposition 6.5). In particular, the open submanifold $\mu^{-1}(\mu(M)^{\text{int}})$, with the restricted symplectic structure, is symplectomorphic to $\mu(M)^{\text{int}} \times T$ with the symplectic structure ω such that $p_1 : \mu(M)^{\text{int}} \times T \rightarrow \mathfrak{t}^*$ is a momentum map for the torus action $t \cdot (\xi, s) = (\xi, ts)$. Thus, to obtain a lower bound on the Gromov width of M , it is sufficient to obtain a lower bound on the Gromov width of $\mu(M)^{\text{int}} \times T$. A lower bound on the Gromov width of $\mu(M)^{\text{int}} \times T$ can be extracted from the combinatorics of $\mu(M)$, as we will now explain. Define

$$\Delta^n(l) = \{(y_1, \dots, y_n) \in \mathbb{R}^n : 0 < y_1, \dots, y_n, \text{ and } y_1 + \dots + y_n < l\},$$

the standard open simplex of size l . A version of the following Lemma was proven in [19] for $n = 2$. Proofs for arbitrary n can be found in [16, 17].

Recall that if (V_i, L_i) , $i = 1, 2$, are vector spaces V_i together with full rank lattices L_i , then an *integral affine map* $B : V_1 \rightarrow V_2$ is a map of the form $B = A + v$ where $A : V_1 \rightarrow V_2$ is a linear map that sends a \mathbb{Z} -basis for L_1 to a \mathbb{Z} -basis for L_2 and $v \in V_2$.

Lemma 12. *Let T be a compact torus of dimension n and let S be an open subset of \mathfrak{t}^* . If there is an integral affine map $B : (\mathbb{R}^n, \mathbb{Z}^n) \rightarrow (\mathfrak{t}^*, L^*)$ such that $B(\Delta^n(l)) \subset S$, then*

$$l \leq G\text{Width}(S \times T, \omega).$$

If a line segment in (\mathbb{R}^n, L) can be translated to a scalar multiple of a primitive vector in the lattice L , then that scalar is the *integral affine length* of the line segment. A vertex v of a convex polytope $\mathcal{P} \subset \mathbb{R}^n$ is *smooth* if the edges of \mathcal{P} incident to v are spanned by a \mathbb{Z} -basis for L (after translation by v). If v is a smooth vertex and the minimum integral affine length of the edges incident to v is l , then there exists an integral affine map B such that $B(\Delta^n(l)) \subset \mathcal{P}^{\text{int}}$.

The other half of Theorem 1 follows from

Theorem 13. [4] *If \mathcal{O}_λ is the G_2 coadjoint orbit through $(0, \lambda)$ as in Figure 1, then*

$$G\text{Width}(\mathcal{O}_\lambda, \omega_\lambda) \leq \frac{\lambda}{\sqrt{3}}.$$

We are now equipped to prove the main theorem.

of Theorem 1.1. Recall that the pre-image of \mathcal{T}^{int} under $\Lambda \circ \Phi$ is a connected open dense submanifold $U \subset \mathcal{O}_\lambda$. The restriction

$$\Lambda \circ \Phi : U \rightarrow \mathcal{C} = \mathcal{T}^{\text{int}}$$

is a proper map to an open convex set that generates a completely integrable Hamiltonian torus action. Thus by [10], (U, ω_λ) is the unique non-compact toric manifold with moment polytope $\Lambda \circ \Phi(U) = \Delta_\lambda \cap \mathcal{C} \subset (\mathbb{R}^5, L^*)$.

It is straightforward to check from Table 1 that the minimum integral affine length of an edge of Δ_λ (with respect to the weight lattice L^* in Equation 8) is $\frac{\lambda}{\sqrt{3}}$. Any vertex of Δ_λ contained in \mathcal{C} (of which there are several) is smooth, so by the discussion following Lemma

12 there exists an integral affine map $B : (\mathbb{R}^5, \mathbb{Z}^5) \rightarrow (\mathbb{R}^5, L^*)$ that embeds $\Delta^5\left(\frac{\lambda}{\sqrt{3}}\right)$ into $\Delta_\lambda^{\text{int}}$. Thus by Lemma 12 and the discussion preceding it we have shown that

$$\frac{\lambda}{\sqrt{3}} \leq \text{GWidth}(\Delta_\lambda^{\text{int}} \times T, \omega) \leq \text{GWidth}(\mathcal{O}_\lambda, \omega_\lambda).$$

Combining this with the upper bound of [4] completes the proof. \square

REFERENCES

- [1] AGRICOLA, I. Old and new on the exceptional group G_2 . *Notices of the American Mathematical Society* 55, 8 (2008), 922–929.
- [2] BIRTEA, P., ORTEGA, J.-P., AND RATIU, T. S. Openness and convexity for momentum maps. *Transactions of the American Mathematical Society* 2, 361 (2009), 603–630.
- [3] BJORND AHL, C., AND KARSHON, Y. Revisiting Tietze-Nakajima: local and global convexity for maps. *Canad. J. Math.* 62, 5 (2010), 975–993.
- [4] CAVIEDES CASTRO, A. Upper bound for the Gromov width of coadjoint orbits of compact Lie groups. Pre-print. arXiv:1404.4647, April 2014.
- [5] GUILLEMIN, V., AND PRATO, E. Heckman, Kostant, and Steinberg formulas for symplectic manifolds. *Advances in Mathematics* 82 (1990), 160–179.
- [6] GUILLEMIN, V., AND STERNBERG, S. The Gelfand-Cetlin system and quantization of the complex flag manifolds. *J. Funct. Anal.* 52, 1 (1983), 106–128.
- [7] GUILLEMIN, V., AND STERNBERG, S. On collective complete integrability according to the method of Thimm. *Ergodic Theory Dynam. Systems* 3, 2 (1983), 219–230.
- [8] HALACHEVA, I., AND PABINIAK, M. The Gromov width of coadjoint orbits of the symplectic group. Pre-print. arXiv:1601.02825, January 2016.
- [9] HARADA, M., AND KAVEH, K. Integrable systems, toric degenerations and Okounkov bodies. *Inventiones mathematicae* 202, 3 (2015), 927–985.
- [10] KARSHON, Y., AND LERMAN, E. Non-compact symplectic toric manifolds. *Symmetry, Integrability and Geometry: Methods and Applications* 11, 055 (2015), 37.
- [11] KARSHON, Y., AND TOLMAN, S. The Gromov width of complex Grassmannians. *Geometry and Topology* 5 (2005), 911–922.
- [12] KAVEH, K. Toric degenerations and symplectic geometry of smooth projective varieties. Pre-print. arXiv:1508.00316, August 2015.
- [13] LANE, J. Convexity and thimm’s trick. Pre-print. arXiv:1509.07356, November 2015.
- [14] LOI, A., AND ZUDDAS, F. On the Gromov width of homogeneous Kaehler manifolds. Pre-print. arXiv:1508.02862, August 2015.
- [15] LU, G. Gromov-Witten invariants and pseudo symplectic capacities. *Israel Journal of Mathematics* 156 (2006), 1–63.
- [16] PABINIAK, M. Gromov width of non-regular coadjoint orbits of $U(n)$, $SO(2n)$ and $SO(2n+1)$. *Mathematical Research Letters* 21, 1 (2014), 187–205.
- [17] SCHLENK, F. On symplectic folding. Pre-print. arXiv:math/9903086, March 1999.
- [18] THIMM, A. Integrable geodesic flows on homogeneous spaces. *Ergodic Theory Dynamical Systems* 1, 4 (1981).
- [19] TRAYNOR, L. Symplectic packing obstructions. *Journal of Differential Geometry* 42, 2 (September 1995).
- [20] WOODWARD, C. The classification of transversal multiplicity-free group actions. *Annals of Global Analysis and Geometry* 14 (1996), 3–42.
- [21] ZOGHI, M. *The Gromov width of coadjoint orbits of the symplectic group Orbits of Compact Lie Groups*. PhD thesis, University of Toronto, 2010.

DEPT. OF MATHEMATICS, UNIVERSITY OF TORONTO, 40 ST. GEORGE STREET, TORONTO ONTARIO M5S 2E4, CANADA

E-mail address: lanejere@math.toronto.edu